arXiv:hep-lat/0109030v1 26 Sep 2001

External Field Dependence of Deconfinement Temperature in SU(3)

Paolo Cea[a] and Leonardo Cosmai[b]

[a]Dipartimento di Fisica, Univ. of Bari and INFN - Sezione di Bari, I-70126 Bari, Italy

[b]INFN - Sezione di Bari, I-70126 Bari, Italy

We study vacuum dynamics of SU(3) lattice gauge theory at finite temperature. Using the lattice Schrödinger functional, SU(3) vacuum is probed by means of an external constant Abelian chromomagnetic field. Our preliminary numerical data suggest that, by increasing the strength of the applied external field, deconfinement temperature decreases towards zero. This means that strong enough Abelian chromomagnetic fields destroy confinement of color.

To study vacuum structure of lattice gauge theories we introduced [1,2] a gauge invariant effective action, defined by means of the lattice Schrödinger functional

$$\mathcal{Z}[U^{\text{ext}}_\mu] = \int \mathcal{D}U \; e^{-S_W} \,. \tag{1}$$

S_W is the standard Wilson action and the functional integration is extended over links on a lattice $L_s^3 \times L_t$ with the hypertorus geometry and satisfying the constraints (x_t: temporal coordinate)

$$U_k(x)|_{x_t=0} = U^{\text{ext}}_k(x) \,, \quad (k=1,2,3) \,, \tag{2}$$

$U^{\text{ext}}_k(x)$ being the lattice version of the external continuum gauge field $\vec{A}^{\text{ext}}(x) = \vec{A}^{\text{ext}}_a(x)\lambda_a/2$.

The lattice effective action for the external static background field $\vec{A}^{\text{ext}}(x)$ is given by

$$\Gamma[\vec{A}^{\text{ext}}] = -\frac{1}{L_4} \ln \left\{ \frac{\mathcal{Z}[\vec{A}^{\text{ext}}]}{\mathcal{Z}[0]} \right\} \tag{3}$$

and is invariant for gauge transformations of the external links U^{ext}_k.

At finite temperature $T = 1/aL_t$, we introduced the thermal partition function in presence of a given static background field:

$$\mathcal{Z}_T \left[\vec{A}^{\text{ext}}\right] = \int_{U_k(L_t,\vec{x})=U_k(0,\vec{x})=U^{\text{ext}}_k(\vec{x})} \mathcal{D}U \, e^{-S_W} \,. \tag{4}$$

In this case the relevant quantity is the free energy functional

$$F[\vec{A}^{\text{ext}}] = -\frac{1}{L_t} \ln \frac{\mathcal{Z}_T[\vec{A}^{\text{ext}}]}{\mathcal{Z}_T[0]} \,. \tag{5}$$

If physical temperature is sent to zero the thermal functional Eq. (4) reduces to Eq. (1). We used the above defined lattice effective action to investigate if deconfinement temperature depends on the strength of an applied external constant Abelian chromomagnetic field.

The static constant Abelian chromomagnetic field in the continuum is

$$\vec{A}^{\text{ext}}_a(\vec{x}) = \vec{A}^{\text{ext}}(\vec{x})\delta_{a,3} \,, \quad A^{\text{ext}}_k(\vec{x}) = \delta_{k,2} x_1 H \,. \tag{6}$$

On the lattice

$$\begin{aligned} &U^{\text{ext}}_1(\vec{x}) = U^{\text{ext}}_3(\vec{x}) = \mathbf{1} \,, \\ &U^{\text{ext}}_2(\vec{x}) = \begin{bmatrix} \exp(i\frac{gHx_1}{2}) & 0 & 0 \\ 0 & \exp(-i\frac{gHx_1}{2}) & 0 \\ 0 & 0 & 1 \end{bmatrix} \end{aligned} \tag{7}$$

Due to periodic boundary conditions the magnetic field H turns out to be quantized

$$\frac{a^2 gH}{2} = \frac{2\pi}{L_1} n_{\text{ext}} \,, \qquad n_{\text{ext}} \ \text{integer} \,. \tag{8}$$

This field gives rise to constant field strength, then due to gauge invariance it is easy to show that $F[\vec{A}^{\text{ext}}]$ is proportional to spatial volume

$V = L_s^3$. Therefore the relevant quantity is the density $f[\vec{A}^{\rm ext}]$ of free energy

$$f[\vec{A}^{\rm ext}] = \frac{1}{V} F[\vec{A}^{\rm ext}] \,. \tag{9}$$

The β-derivative of $f[\vec{A}^{\rm ext}]$ (at fixed external field strength gH) is easily evaluated ($\Omega = L_s^3 \times L_t$)

$$f'[\vec{A}^{\rm ext}] = \left\langle \frac{1}{\Omega} \sum_{x \in \tilde{\Lambda}, \mu < \nu} \frac{1}{3} {\rm Re Tr} U_{\mu\nu}(x) \right\rangle_0 - \left\langle \frac{1}{\Omega} \sum_{x \in \tilde{\Lambda}, \mu < \nu} \frac{1}{3} {\rm Re Tr} U_{\mu\nu}(x) \right\rangle_{\vec{A}^{\rm ext}} \,. \tag{10}$$

The generic plaquette $U_{\mu\nu}(x) = U_\mu(x) U_\nu(x + \hat{\mu}) U_\mu^\dagger(x + \hat{\nu}) U_\nu^\dagger(x)$ contributes to the sum in Eq. (10) if the link $U_\mu(x)$ is a "dynamical" one (i.e. it is not constrained in the functional integration Eq. (4)). $\tilde{\Lambda}$ is a sub-ensemble of the lattice sites, $x \in \tilde{\Lambda}$ if the link $U_\mu(x)$ exiting from it is a dynamical link.

As is well known pure SU(3) gauge system undergoes a deconfinement phase transition by increasing temperature. To estimate deconfinement temperature T_c, we evaluate $f'[\vec{A}^{\rm ext}]$ as a function of β for different lattice temporal sizes L_t. To determine the pseudocritical couplings we parameterize $f'(\beta, L_t)$ (see Fig. 1) near the peak as

$$\frac{f'(\beta, L_t)}{\varepsilon'_{\rm ext}} = \frac{a_1(L_t)}{a_2(L_t)[\beta - \beta^*(L_t)]^2 + 1} \,. \tag{11}$$

In Eq. (11) $\varepsilon'_{\rm ext}$ is the derivative of the classical energy due to the external applied field

$$\varepsilon'_{\rm ext} = \frac{2}{3} \left[1 - \cos(\frac{gH}{2})\right] = \frac{2}{3} \left[1 - \cos(\frac{2\pi}{L_1} n_{\rm ext})\right]. \tag{12}$$

Once determined $\beta^*(L_t)$ we estimate the deconfinement temperature as

$$\frac{T_c}{\Lambda_{\rm latt}} = \frac{1}{L_t} \frac{1}{f(\beta^*(L_t))} \,, \tag{13}$$

with

$$f(\beta) = \left(\frac{\beta}{2 N b_0} \right)^{b_1/2b_0^2} \exp \left(-\beta \frac{1}{4 N b_0} \right) \,, \tag{14}$$



Figure 1. f' versus β for various L_t at given external field strength ($n_{\rm ext} = 1$.)

where N is the color number, $b_0 = (11N)/(48\pi^2)$, $b_1 = (34N^2)/(3(16\pi^2)^2)$.

Following [3] we perform a linear extrapolation to the continuum of our data for $T_c/\Lambda_{\rm latt}$. We see that our estimate of $T_c/\Lambda_{\rm latt}$ in the continuum is in fair agreement with the one available in the literature [3] without external field (see Fig. 2). However Fig. 2 is suggestive of a possible non-trivial dependence of deconfinement critical temperature on external Abelian chromomagnetic field. Therefore we decided to vary the strength of the applied external Abelian chromomagnetic background field to study possible dependence of T_c on gH. To this aim we performed numerical simulations on $64^3 \times L_t$ lattices with $n_{\rm ext} = 1, 5, 10$. In Fig. 3 we display our determination of T_c for two different values ($n_{\rm ext} = 1, 10$) of the applied field strength. We see that the critical temperature decreases by increasing the external Abelian chromomagnetic field. For dimensional reasons one expects that

$$T_c^2 \;\sim\; gH \,. \tag{15}$$

Indeed we try to fit our data with

$$\frac{T_c(gH)}{\Lambda_{\rm latt}} = \frac{T_c(0)}{\Lambda_{\rm latt}} + \alpha \sqrt{gH} \,. \tag{16}$$

We find a satisfying fit with $\alpha = -27.6 \pm 2.6$ (see



Figure 2. The estimate of T_c for $n_{\rm ext} = 1$ and $n_{\rm ext} = 0$.



Figure 3. The estimate of T_c for $n_{\rm ext} = 1$ and $n_{\rm ext} = 10$.

Fig. 4). Remarkably, we see that there exists a critical field

$$a^2 gH_c \simeq 1.15 \tag{17}$$

such that $T_c = 0$ for $H > H_c$.



Figure 4. $T_c/\Lambda_{\rm LAT}$ versus the applied external field strength gH.

In conclusion, we found [2] that at zero-temperature, in the thermodynamic limit, SU(3) vacuum screens completely external chromomagnetic Abelian field. Moreover, by increasing temperature there is no screening of the Abelian chromomagnetic field. There is a critical field H_c such that, for $H > H_c$, the gauge system is in the deconfined phase. In particular, this last result suggests that there is an intimate connection between Abelian chromomagnetic fields and color confinement.

REFERENCES

1. P. Cea, L. Cosmai and A.D. Polosa, Phys. Lett. B392 (1997) 177, hep-lat/9601010.
2. P. Cea and L. Cosmai, Phys. Rev. D60 (1999) 094506, hep-lat/9903005.
3. J. Fingberg, U. Heller and F. Karsch, Nucl. Phys. B392 (1993) 493, hep-lat/9208012.